

MATH


QA

372

.081

1886a

MATH
QA
372
081
1886a



CORNELL UNIVERSITY LIBRARY


3 1924 067 741 995

GAYLORD

PRINTED IN U.S.A.

GAYLORD

PRINTED IN U.S.A



Cornell University
Library


The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

<http://www.archive.org/details/cu31924067741995>

Production Note

Cornell University Library produced this volume to replace the irreparably deteriorated original. It was scanned using Xerox software and equipment at 600 dots per inch resolution and compressed prior to storage using CCITT Group 4 compression. The digital data were used to create Cornell's replacement volume on paper that meets the ANSI Standard Z39.48-1984. The production of this volume was supported in part by the Commission on Preservation and Access and the Xerox Corporation. Digital file copyright by Cornell University Library 1992.



Cornell University Library
Ithaca, New York

FROM

John Henry Tanner.....

.....
MATHEMATICS
.....

EXAMPLES
OF
DIFFERENTIAL EQUATIONS

WITH
RULES FOR THEIR SOLUTION.

BY
GEORGE A. OSBORNE, S.B.
PROFESSOR OF MATHEMATICS IN THE MASSACHUSETTS INSTITUTE
OF TECHNOLOGY.



BOSTON, U.S.A. :
PUBLISHED BY GINN & COMPANY.
1889.



Entered, according to Act of Congress, in the year 1886, by
GEORGE A. OSBORNE,
in the Office of the Librarian of Congress, at Washington.

PREFACE.

THIS work has been prepared to meet a want felt by the author in a practical course on the subject, arranged for advanced students in Physics. It is intended to be used in connection with lectures on the theory of Differential Equations and the derivation of the methods of solution.

Many of the examples have been collected from standard treatises, but a considerable number have been prepared by the author to illustrate special difficulties, or to provide exercises corresponding more nearly with the abilities of average students. With few exceptions they have all been tested by use in the class-room.

G. A. OSBORNE.

Boston, Feb. 1, 1886.

CONTENTS.

CHAPTER I.

DEFINITIONS. — DERIVATION OF THE DIFFERENTIAL EQUATION FROM THE COMPLETE PRIMITIVE.

	PAGE.
Definitions	1
Derivation of differential equations of the first order	1
Derivation of differential equations of higher orders	2

SOLUTION OF DIFFERENTIAL EQUATIONS.

CHAPTER II.

DIFFERENTIAL EQUATIONS OF FIRST ORDER AND DEGREE BETWEEN TWO VARIABLES.

Form, $XYdx + X'Y'dy = 0$	4
Homogeneous equations	5
Form, $(ax + by + c)dx + (a'x + b'y + c')dy = 0$	5
Linear form, $\frac{dy}{dx} + Py = Q$	6
Form, $\frac{dy}{dx} + Py = Qy^n$	7

CHAPTER III.

EXACT DIFFERENTIAL EQUATIONS. — INTEGRATING FACTORS.

Solution of exact differential equations	8
--	---

	PAGE.
Solution by means of an integrating factor in the following cases :	
Homogeneous equations	9
Form, $f_1(xy)y dx + f_2(xy)x dy = 0$	10
When $\frac{\frac{dM}{dy} - \frac{dN}{dx}}{N} = \phi(x)$, or $\frac{\frac{dN}{dx} - \frac{dM}{dy}}{M} = \psi(y)$	10

CHAPTER IV.

DIFFERENTIAL EQUATIONS OF FIRST ORDER AND DEGREE WITH THREE
VARIABLES.

Condition for a single primitive, and method of solution	12
--	----

CHAPTER V.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER OF HIGHER DEGREES.

Equations which can be solved with respect to p	14
Equations which can be solved with respect to y	14
Equations which can be solved with respect to x	15
Homogeneous equations	16
Clairaut's form, $y = px + f(p)$	16

CHAPTER VI.

SINGULAR SOLUTIONS.

Method of deriving the singular solution either from the complete primitive or from the differential equation	17
--	----

DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

CHAPTER VII.

LINEAR DIFFERENTIAL EQUATIONS.

Linear equations with constant coefficients and second member zero ..	19
Linear equations with constant coefficients and second member not zero,	22
A special form of linear equations with variable coefficients	24

CHAPTER VIII.

SPECIAL FORMS OF DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

	PAGE.
Form, $\frac{d^n y}{dx^n} = X$	25
Form, $\frac{d^2 y}{dx^2} = Y$	25
Equations not containing y directly	26
Equations not containing x directly	27

CHAPTER IX.

SIMULTANEOUS DIFFERENTIAL EQUATIONS.

Simultaneous equations of first order	28
Simultaneous equations of higher orders	30

CHAPTER X.

Geometrical applications	32
--------------------------------	----

ANSWERS TO EXAMPLES	35
---------------------------	----

EXAMPLES OF DIFFERENTIAL EQUATIONS.



CHAPTER I.

DEFINITIONS. DERIVATION OF THE DIFFERENTIAL EQUATION FROM THE COMPLETE PRIMITIVE.

1. A *differential equation* is an equation containing differentials or differential coefficients.

The *solution* of a differential equation is the determination of another equation free from differentials or differential coefficients, from which the former may be derived by differentiation.

The *order* of a differential equation is that of the highest differential coefficient it contains; and its *degree* is that of the highest power to which this highest differential coefficient is raised, after the equation is freed from fractions and radicals.

The solution of a differential equation requires one or more integrations, each of which introduces an arbitrary constant. The most general solution of a differential equation of the n th order contains n arbitrary constants, whatever may be its degree. This general solution is called the *complete primitive* of the given differential equation.

2. To derive a differential equation of the first order from its complete primitive.

Differentiate the primitive; and if the arbitrary constant has disappeared, the result is the required differential equation. If not, the elimination of this constant between the two equations will give the differential equation.

2 DERIVATION OF THE DIFFERENTIAL EQUATION.

3. Form the differential equations of the first order of which the following are the complete primitives, c being the arbitrary constant :

1. $\log(xy) + x = y + c.$
2. $(1 + x^2)(1 + y^2) = cx^2.$
3. $\cos y = c \cos x.$
4. $y = ce^{-\tan^{-1}x} + \tan^{-1}x - 1.$
5. $y = (cx + \log x + 1)^{-1}.$
6. $y = cx + c - c^3.$
7. $(y + c)^2 = 4ax.$
8. $y^2 \sin^2 x + 2cy + c^2 = 0.$
9. $e^{2y} + 2cxe^y + c^2 = 0.$

4. *To derive a differential equation of the second order from its complete primitive.*

Differentiate the primitive twice successively, and eliminate, if necessary, the two arbitrary constants between the three equations.

5. Form the differential equations of the second order of which the following are the complete primitives, c_1 and c_2 being the arbitrary constants :

1. $y = c_1 \cos(ax + c_2).$
2. $y = c_1 e^{ax} + c_2 e^{-ax}.$
3. $y = (c_1 + c_2 x) e^{ax}.$
4. $y = c_1 x^3 + \frac{c_2}{x}.$
5. $y = c_1 \sin nx + c_2 \cos nx + \frac{\cos ax}{n^2 - a^2}.$

6. The preceding process may be extended to the derivation of equations of higher orders from their primitives.

7. Form the differential equations of the third order of which the following are the complete primitives :

1. $y = c_1 e^{2x} + c_2 e^{-3x} + c_3 e^x.$

2. $ye^x = c_1 e^{2x} + c_2 \sin x \sqrt{2} + c_3 \cos x \sqrt{2}.$

3. $y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x + c_3.$

Form the differential equations of the fourth order of which the following are the complete primitives :

4. $y = (c_1 + c_2 x + c_3 x^2) e^x + c_4.$

5. $x^3 + a^4 y = c_1 e^{ax} + c_2 e^{-ax} + c_3 \sin ax + c_4 \cos ax.$

CHAPTER II.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND FIRST DEGREE BETWEEN TWO VARIABLES.

General Form, $Mdx + Ndy = 0$,

where M, N , are each functions of x and y .

8. Form, $XYdx + X'Y'dy = 0$,

where X, X' , are functions of x alone, and Y, Y' , functions of y alone.

Divide so as to separate the variables, and integrate each part separately.

9. Solve the following equations :

1. $(1+x)ydx + (1-y)xdy = 0$.

2. $(x^2 - yx^2) \frac{dy}{dx} + y^2 + xy^2 = 0$.

3. $\frac{dy}{dx} = \frac{1+y^2}{(1+x^2)xy}$.

4. $a \left(x \frac{dy}{dx} + 2y \right) = xy \frac{dy}{dx}$.

5. $(1+y^2)dx = (y + \sqrt{1+y^2})(1+x^2)^{\frac{3}{2}}dy$.

6. $\sin x \cos y dx = \cos x \sin y dy$.

7. $\sec^2 x \tan y dx + \sec^2 y \tan x dy = 0$.

8. $\sec^2 x \tan y dy + \sec^2 y \tan x dx = 0$.

9. $\frac{dy}{dx} + \frac{1+y+y^2}{1+x+x^2} = 0$.

10. *Homogeneous equations.*

Substitute $y = vx$; in the resulting equation between v and x , the variables can be separated. (See Art. 8.)

11. Solve the following equations :

$$1. \quad (y - x)dy + ydx = 0.$$

$$2. \quad (2\sqrt{xy} - x)dy + ydx = 0.$$

$$3. \quad y^2 + x^2 \frac{dy}{dx} = xy \frac{dy}{dx}.$$

$$4. \quad x \frac{dy}{dx} = y + \sqrt{x^2 + y^2}.$$

$$5. \quad x \cos \frac{y}{x} \cdot \frac{dy}{dx} = y \cos \frac{y}{x} - x.$$

$$6. \quad (8y + 10x)dx + (5y + 7x)dy = 0.$$

$$7. \quad (x + y) \frac{dy}{dx} = y - x.$$

$$8. \quad x \cos \frac{y}{x} (y dx + x dy) = y \sin \frac{y}{x} (x dy - y dx).$$

$$9. \quad x + y \frac{dy}{dx} = my.$$

$$(1), \quad m < 2; \quad (2), \quad m = 2; \quad (3), \quad m > 2.$$

$$10. \quad [(x^2 - y^2) \sin a + 2xy \cos a - y \sqrt{x^2 + y^2}] \frac{dy}{dx} \\ = 2xy \sin a - (x^2 - y^2) \cos a + x \sqrt{x^2 + y^2}.$$

12. Form,

$$(ax + by + c)dx + (a'x + b'y + c')dy = 0.$$

Substitute $x = x' + a, \quad y = y' + \beta,$

and determine the constants a, β , so that the new equation between x' and y' may be homogeneous. (See Art. 10.)

This method fails when $\frac{a}{a'} = \frac{b}{b'}$. In this case put $ax + by = z$, and obtain a new equation between x and z or between y and z ; the variables can then be separated.

13. Solve the following equations :

$$1. \quad (3y - 7x + 7)dx + (7y - 3x + 3)dy = 0.$$

$$2. \quad (4x + 2y - 1) \frac{dy}{dx} + 2x + y + 1 = 0.$$

$$3. \quad \frac{dy}{dx} = \frac{7y + x + 2}{3x + 5y + 6}.$$

$$4. \quad (2y + x + 1)dx = (2x + 4y + 3)dy.$$

$$5. \quad 2x - y + 1 + (x + y - 2) \frac{dy}{dx} = 0.$$

14. *Linear Form*, $\frac{dy}{dx} + Py = Q$,

where P, Q , are independent of y .

Solution,
$$y = e^{-\int P dx} \left(\int Q e^{\int P dx} dx + c \right).$$

15. Solve the following equations :

$$1. \quad x \frac{dy}{dx} - ay = x + 1.$$

$$2. \quad x(1 - x^2)dy + (2x^2 - 1)ydx = ax^3dx.$$

$$3. \quad (1 - x^2)^2 \frac{dy}{dx} + y\sqrt{1 - x^2} = x + \sqrt{1 - x^2}.$$

$$4. \quad \frac{dy}{dx} + y \cos x = \frac{1}{2} \sin 2x.$$

$$5. \quad (1 + y^2)dx = (\tan^{-1}y - x) dy.$$

$$6. \quad \sqrt{a^2 + x^2} \left(1 - \frac{dy}{dx} \right) = x + y.$$

$$7. \quad (1 + x^2) dy + \left(xy - \frac{1}{x} \right) dx = 0.$$

$$8. \quad \frac{dy}{dx} + y \frac{d\phi}{dx} = \phi \frac{d\phi}{dx},$$

where ϕ is a function of x alone.

$$16. \text{ Form, } \frac{dy}{dx} + Py = Qy^n,$$

where P, Q , are independent of y .

Divide by y^n , and substitute $z = y^{-n+1}$. The new equation between z and x will be linear. (See Art. 14.)

17. Solve the following equations :

$$1. \quad (1 - x^2) \frac{dy}{dx} - xy = axy^2.$$

$$2. \quad 3y^2 \frac{dy}{dx} - ay^3 = x + 1.$$

$$3. \quad \frac{dy}{dx} = 2xy (ax^2y^2 - 1).$$

$$4. \quad \frac{dy}{dx} (x^2y^3 + xy) = 1.$$

$$5. \quad \frac{dy}{dx} + y \cos x = y^n \sin 2x.$$

$$6. \quad (y \log x - 1) y dx = x dy.$$

$$7. \quad ax^2y^n dy + y dx = 2x dy.$$

$$8. \quad y - \cos x \frac{dy}{dx} = y^2 \cos x (1 - \sin x).$$

$$9. \quad y \frac{dy}{dx} + by^2 = a \cos x.$$

CHAPTER III.

EXACT DIFFERENTIAL EQUATIONS AND INTEGRATING FACTORS.

18. $Mdx + Ndy$ is an exact differential when

$$\frac{dM}{dy} = \frac{dN}{dx}. \quad (1)$$

The solution of

$$Mdx + Ndy = 0, \quad \text{in this case is}$$

$$\int Mdx + \int \left(N - \frac{d}{dy} \int Mdx \right) dy = c,$$

or
$$\int Ndy + \int \left(M - \frac{d}{dx} \int Ndy \right) dx = c.$$

In integrating with respect to x , y is regarded as constant, and in integrating with respect to y , x is regarded as constant.

19. Solve the following equations after applying the condition (1) for an exact differential:

1. $(x^3 + 3xy^2)dx + (y^3 + 3x^2y)dy = 0.$

2. $(x^2 - 4xy - 2y^2)dx + (y^2 - 4xy - 2x^2)dy = 0.$

3. $\left(1 + \frac{y^2}{x^2}\right)dx - \frac{2y}{x}dy = 0.$

4. $\frac{2x dx}{y^3} + \left(\frac{1}{y^2} - \frac{3x^2}{y^4}\right)dy = 0.$

5. $x dx + y dy + \frac{x dy - y dx}{x^2 + y^2} = 0.$
6. $\frac{dx}{\sqrt{x^2 + y^2}} + \left(1 - \frac{x}{\sqrt{x^2 + y^2}}\right) \frac{dy}{y} = 0.$
7. $\left(x + \frac{1}{\sqrt{y^2 - x^2}}\right) dx + \left(y - \frac{x}{y \sqrt{y^2 - x^2}}\right) dy = 0.$
8. $(1 + e^{\frac{x}{y}}) dx + e^{\frac{x}{y}} \left(1 - \frac{x}{y}\right) dy = 0.$
9. $e^x (x^2 + y^2 + 2x) dx + 2y e^x dy = 0.$
10. $(m dx + n dy) \sin(mx + ny) = (n dx + m dy) \cos(nx + my).$
11. $\frac{x dx + y dy}{\sqrt{1 + x^2 + y^2}} + \frac{y dx - x dy}{x^2 + y^2} = 0.$
12. $\frac{x^a dy - a y x^{a-1} dx}{b y^2 - g x^{2a}} = x^{a-1} dx.$
 (1), $g > 0$; (2), $g < 0$ and $= -k$; (3), $g = 0$;
 (4), $a = 0$; (5), $b = 0.$

20. When $Mdx + Ndy$ is not an exact differential, it may sometimes be made exact by multiplying by a factor, called an *integrating factor*. The following are some of the cases where this is possible.

21. When $Mdx + Ndy$ is homogeneous, $\frac{1}{Mx + Ny}$ is an integrating factor. This fails when $Mx + Ny = 0$, but in that case the solution is $y = cx$.

22. Solve the following equations by means of an integrating factor :

1. $(x^2 + 2xy - y^2) dx = (x^2 - 2xy - y^2) dy.$
2. $\frac{dx}{x} + \frac{dy}{y} + 2\left(\frac{dx}{y} - \frac{dy}{x}\right) = 0.$

$$3. \quad (x^2 y^2 + xy^3) dx - (x^3 y + x^2 y^2) dy = 0.$$

$$4. \quad x^3 dx + (3x^2 y + 2y^3) dy = 0.$$

$$5. \quad (x\sqrt{x^2 + y^2} - x^2) dy + (xy - y\sqrt{x^2 + y^2}) dx = 0.$$

(See Art. 11 for other examples.)

$$23. \text{ Form, } f_1(xy)y dx + f_2(xy)x dy = 0.$$

$\frac{1}{Mx - Ny}$ is an integrating factor. This fails when

$$Mx - Ny = 0,$$

but in that case the solution is $xy = c$.

Another method of solving is to put $xy = v$, and obtain an equation between x and v or between y and v . The variables can then be separated.

24. Solve the following equations by means of an integrating factor :

$$1. \quad (1 + xy)y dx + (1 - xy)x dy = 0.$$

$$2. \quad (x^2 y^2 + xy)y dx + (x^2 y^2 - 1)x dy = 0.$$

$$3. \quad (x^3 y^3 + 1)(x dy + y dx) + (x^2 y^2 + xy)(y dx - x dy) = 0.$$

$$4. \quad (\sqrt{xy} - 1)x dy - (\sqrt{xy} + 1)y dx = 0.$$

$$5. \quad (y + y\sqrt{xy})dx + (x + x\sqrt{xy})dy = 0.$$

$$6. \quad e^{xy}(x^2 y^2 + xy)(x dy + y dx) + y dx - x dy = 0.$$

$$7. \quad xy[1 + \cot(xy)](x dy + y dx) + x dy - y dx = 0.$$

$$25. \text{ When } \frac{\frac{dM}{dy} - \frac{dN}{dx}}{N} = \phi(x),$$

then $e^{\int \phi(x) dx}$ is an integrating factor.

$$\text{Or, when } \frac{\frac{dN}{dx} - \frac{dM}{dy}}{M} = \psi(y),$$

then $e^{\int \psi(y) dy}$ is an integrating factor.

26. Solve the following equations by means of an integrating factor :

1. $(x^2 + y^2 + 2x)dx + 2ydy = 0.$

2. $(3x^2 - y^2) \frac{dy}{dx} = 2xy.$

3. $\frac{dy}{dx} = \frac{x^2 + y^2}{2xy}.$

4. $[(1-y)\sqrt{1-x^2} - xy]dx + [1-x^2 - x\sqrt{1-x^2}]dy = 0.$

5. $(\cos x + 2y \sec x \sec^2 2x)dx + (\tan 2x \sec y - \sin x \tan y)dy = 0.$

6. $\sin(3x - 2y)(2dx - dy) + \sin(x - 2y)dy = 0.$

7. The Linear Equation

$$\frac{dy}{dx} + Py = Q,$$

where P and Q are independent of y .

CHAPTER IV.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND DEGREE CONTAINING THREE VARIABLES.

General form, $Pdx + Qdy + Rdz = 0$,

where P, Q, R , are each functions of x, y, z .

27. If the variables can be separated, solve by integrating the parts separately.

The equation is derivable from a single primitive only when the following condition is satisfied :

$$P\left(\frac{dQ}{dz} - \frac{dR}{dy}\right) + Q\left(\frac{dR}{dx} - \frac{dP}{dz}\right) + R\left(\frac{dP}{dy} - \frac{dQ}{dx}\right) = 0. \quad (1)$$

The solution may then be effected by first solving the equation with one of the parts Pdx, Qdy, Rdz , omitted, regarding x, y, z , respectively, constant.

Omitting Rdz , for example, we solve $Pdx + Qdy = 0$, regarding z constant, and introducing instead of an arbitrary constant of integration, Z , an undetermined function of z , which must be subsequently determined so that this primitive may satisfy the given differential equation. The equation of condition for determining Z will ultimately involve only Z and z .

28. Solve the following equations after applying the condition (1) for a single primitive :

$$1. \quad \frac{dx}{x-a} + \frac{dy}{y-b} + \frac{dz}{z-c} = 0.$$

$$2. \quad (x-3y-z)dx + (2y-3x)dy + (z-x)dz = 0.$$

3. $(y+z)dx + (z+x)dy + (x+y)dz = 0.$
4. $yz\,dx + zx\,dy + xy\,dz = 0.$
5. $(y+z)dx + dy + dz = 0.$
6. $ay^2z^2\,dx + bz^2x^2\,dy + cx^2y^2\,dz = 0.$
7. $zy\,dx = zx\,dy + y^2\,dz.$
8. $(y\,dx + x\,dy)(a+z) = xy\,dz.$
9. $(y+a)^2\,dx + z\,dy = (y+a)\,dz.$
10. $(y^2 + yz)dx + (xz + z^2)dy + (y^2 - xy)dz = 0.$
11. $(2x^2 + 2xy + 2xz^2 + 1)dx + dy + 2z\,dz = 0.$
12. $(x^2y - y^3 - y^2z)dx + (xy^2 - x^2z - x^3)dy + (xy^2 + x^2y)dz = 0.$

CHAPTER V.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER, OF A DEGREE ABOVE THE FIRST.

In what follows, p denotes $\frac{dy}{dx}$.

29. *When the equation can be solved with respect to p .*

The different values of p constitute so many differential equations of the first degree, which must be solved separately, using the same character for the arbitrary constant in all.

If the terms of each of these separate primitives be transposed to the first member, the product of these first members placed equal to zero will be the complete primitive.

30. Solve the following equations :

1. $p^2 - 5p + 6 = 0.$

2. $x^2 p^2 - a^2 = 0.$

3. $x p^2 - a = 0.$

4. $x p^2 = 1 - x.$

5. $x^2 p^2 + 3 x y p + 2 y^2 = 0.$

6. $p(p + y) = x(x + y).$

7. $p^3 + 2 x p^2 - y^2 p^2 - 2 x y^2 p = 0.$

8. $p^3 - (x^2 + x y + y^2) p^2 + (x^3 y + x^2 y^2 + x y^3) p - x^3 y^3 = 0.$

9. $p^2 + 2 p y \cot x = y^2.$

31. *When the equation can be solved with respect to y .*

Differentiate, regarding p variable as well as x and y , and

substitute for dy , pdx . There will result a differential equation of the first degree between x and p . Solve this equation, and eliminate p between its primitive and the given equation.

32. Solve the following equations :

1. $x - yp = ap^2$.
2. $y = xp^2 + 2p$.
3. $(x + yp)^2 = a^2(1 + p^2)$.
4. $y = xp + p - p^2$.
5. $(y - ap)^2 = 1 + p^2$.
6. $y = ap + bp^2$.
7. $x^2 + y = p^2$.
8. $y^2 = x^2(1 + p^2)$.
9. $y = p^2 + 2p^3$.

33. *When the equation can be solved with respect to x .*

Differentiate, regarding p variable as well as x and y , and substitute for dx , $\frac{dy}{p}$. There will result a differential equation of the first degree between y and p . Solve this equation, and eliminate p between its primitive and the given equation.

34. Solve the following equations :

1. $p^2y + 2px = y$.
2. $x = p + \log p$.
3. $p^2(x^2 + 2ax) = a^2$.
4. $x^2p^2 = 1 + p^2$.
5. $(x - ap)^2 = 1 + p^2$; also when $a = 1$.
6. $x = ap + bp^2$.
7. $my - nxp = yp^2$.

16 HOMOGENEOUS EQUATIONS. — CLAIRAUT'S FORM.

35. When the equation is homogeneous with respect to x and y .

Substitute $y = vx$. If the resulting equation between p and v can be solved with respect to v , the given equation comes under Art. 31 or Art. 33.

But if we can solve with respect to p , substitute for p , $v + x \frac{dv}{dx}$, and there will result a differential equation of the first degree between v and x .

36. Solve the following equations :

1. $xy^2(p^2 + 2) = 2py^3 + x^3$.
2. $(2p + 1)x^{\frac{1}{2}}y = x^{\frac{3}{2}}p^2 + 2y^{\frac{3}{2}}$.
3. $4x^2 = 3(3y - px)(y + px)$.
4. $ds = \left(\frac{y}{2x}\right)^{\frac{1}{2}} dx + \left(\frac{x}{2y}\right)^{\frac{1}{2}} dy$, where $ds = \sqrt{1 + p^2} \cdot dx$.
5. $(nx + py)^2 = (1 + p^2)(y^2 + nx^2)$.

37. Clairaut's Form,

$$y = px + f(p).$$

The solution is immediately obtained by substituting $p = c$.

38. Solve the following equations :

1. $y = px + \frac{m}{p}$.
2. $y = px + p - p^3$.
3. $y^2 - 2pxy - 1 = p^2(1 - x^2)$.
4. $y = 2px + y^2p^3$. Put $y^2 = y'$.
5. $ayp^2 + (2x - b)p = y$. Put $y^2 = y'$.
6. $x^2(y - px) = yp^2$. Put $y^2 = y'$, $x^2 = x'$.
7. $e^{3x}(p - 1) + p^3e^{2y} = 0$. Put $e^x = x'$, $e^y = y'$.
8. $(px - y)(py + x) = h^2p$. Put $y^2 = y'$, $x^2 = x'$.

CHAPTER VI.

SINGULAR SOLUTIONS.

39. A *singular solution* of a differential equation is a solution which is not included in the complete primitive. Differential equations of the first degree have no singular solution. Those of higher degrees may have singular solutions, which may be derived either from the complete primitive, or directly from the differential equation.

40. Let $f(x, y, c) = 0$ be the complete primitive.

By differentiating, regarding c as the only variable, obtain $\frac{df}{dc} = 0$. If we eliminate c between this equation and the primitive, the result will be a singular solution, provided it satisfies the given differential equation.

41. Let $f(x, y, p) = 0$ be the given differential equation.

By differentiating, regarding p as the only variable, obtain $\frac{df}{dp} = 0$. If we eliminate p between this equation and the given differential equation, the result will be a singular solution, provided it satisfies the differential equation.

42. Derive the singular solution of the following equations, directly from the given equation, and also from the complete primitive :

1. $y = px + \frac{m}{p}.$

2. $y^2 - 2xyp + (1 + x^2)p^2 = 1.$

- 3. $p^3 - 4xyp + 8y^2 = 0$. Put $y = z^2$.
- 4. $y = (x - 1)p - p^2$.
- 5. $y(1 + p^2) = 2xp$.
- 6. $x^2p^2 - 2(xy - 2)p + y^2 = 0$.
- 7. $(y - xp)(mp - n) = mnp$.

DIFFERENTIAL EQUATIONS OF AN ORDER HIGHER THAN THE FIRST.



CHAPTER VII.

LINEAR DIFFERENTIAL EQUATIONS.

General Form,

$$\frac{d^n y}{dx^n} + X_1 \frac{d^{n-1} y}{dx^{n-1}} + X_2 \frac{d^{n-2} y}{dx^{n-2}} \dots + X_{n-1} \frac{dy}{dx} + X_n y = X,$$

the coefficients X_1, X_2, \dots, X_n and X being functions of x alone or constants.

43. Linear equations with *constant* coefficients and second member *zero* may be solved as follows :

Substitute in the given equation,

$$\frac{d^n y}{dx^n} = m^n, \quad \frac{d^{n-1} y}{dx^{n-1}} = m^{n-1}, \dots \dots \frac{dy}{dx} = m, \quad y = m^0 = 1.$$

There will result an equation of the n th degree in m , called the *auxiliary equation*. Find the n roots of this equation; these roots will determine a series of terms expressing the complete value of y as follows, viz. For each real root m_1 , there will be a term $Ce^{m_1 x}$; for each pair of imaginary roots $a \pm b\sqrt{-1}$, a term $e^{ax}(A\sin bx + B\cos bx)$; each of the coefficients A, B, C , being an arbitrary constant if the corresponding root occur only once, but a polynomial $c_1 + c_2 x + c_3 x^2 \dots + c_r x^{r-1}$, if the root occur r times.

44. *Roots of auxiliary equation, real and unequal.*

Solve the following equations :

1. $\frac{d^2 y}{dx^2} = a^2 y.$

2. $\frac{d^2 y}{dx^2} + 12y = 7 \frac{dy}{dx}.$

3. $a \frac{d^2 y}{dx^2} = \frac{dy}{dx}.$

4. $3 \left(\frac{d^2 y}{dx^2} + y \right) = 10 \frac{dy}{dx}.$

5. $\frac{d^2 y}{dx^2} + 4 \frac{dy}{dx} = y.$

6. $ab \left(y + \frac{d^2 y}{dx^2} \right) = (a^2 + b^2) \frac{dy}{dx}.$

7. $\frac{d^3 y}{dx^3} = 4 \frac{dy}{dx}.$

8. $\frac{d^3 y}{dx^3} = \frac{d^2 y}{dx^2} + 6 \frac{dy}{dx}.$

9. $\frac{d^3 y}{dx^3} = 7 \frac{dy}{dx} - 6y.$

10. $\frac{d^4 y}{dx^4} + 27y = 12 \frac{d^2 y}{dx^2}.$

11. $\frac{d^5 y}{dx^5} - 2(a^2 + b^2) \frac{d^3 y}{dx^3} + (a^2 - b^2)^2 \frac{dy}{dx} = 0.$

45. *Roots of auxiliary equation unequal, but not all real.*

Solve the following equations :

1. $\frac{d^2 y}{dx^2} + y = 0.$

2. $\frac{d^2 y}{dx^2} - 6 \frac{dy}{dx} + 13y = 0.$

3. $\frac{d^2y}{dx^2} - 2a\frac{dy}{dx} + b^2y = 0.$
(1), when $a > b$; (2), when $a < b$.
4. $\frac{d^2y}{dx^2} - 4ab\frac{dy}{dx} + (a^2 + b^2)^2y = 0.$
5. $\frac{d^2y}{dx^2} - 2\log a\frac{dy}{dx} + \{1 + (\log a)^2\}y = 0.$
6. $\frac{d^3y}{dx^3} + 2\frac{dy}{dx} = 0.$
7. $\frac{d^3y}{dx^3} = y.$
8. $\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} + \frac{dy}{dx} = 3y.$
9. $\frac{d^4y}{dx^4} = y.$
10. $\frac{d^4y}{dx^4} + 2\frac{d^2y}{dx^2} - 8y = 0.$
11. $\frac{d^4y}{dx^4} + 4a^4y = 0.$
12. $\frac{d^6y}{dx^6} = y.$
13. $\frac{d^6y}{dx^6} = -y.$
14. $\frac{d^8y}{dx^8} = y.$

46. *Auxiliary equation containing equal roots.*

Solve the following equations :

1. $\frac{d^2y}{dx^2} - 2a\frac{dy}{dx} + a^2y = 0.$

$$2. \quad \frac{d^2 y}{dx^2} = 0.$$

$$3. \quad \frac{d^3 y}{dx^3} = 4 \frac{d^2 y}{dx^2}.$$

$$4. \quad \frac{d^3 y}{dx^3} - 3 \frac{d^2 y}{dx^2} + 4y = 0.$$

$$5. \quad \frac{d^3 y}{dx^3} - \frac{d^2 y}{dx^2} - \frac{dy}{dx} + y = 0.$$

$$6. \quad \frac{d^4 y}{dx^4} + 2n^2 \frac{d^2 y}{dx^2} + n^4 y = 0.$$

$$7. \quad \frac{d^4 y}{dx^4} - 3 \frac{d^3 y}{dx^3} + 3 \frac{d^2 y}{dx^2} - \frac{dy}{dx} = 0.$$

$$8. \quad \frac{d^4 y}{dx^4} - 4 \frac{d^3 y}{dx^3} + 14 \frac{d^2 y}{dx^2} - 20 \frac{dy}{dx} + 25y = 0, \quad \text{the first member of auxiliary equation being a perfect square.}$$

$$9. \quad \frac{d^n y}{dx^n} + \frac{d^{n-2} y}{dx^{n-2}} = 0.$$

47. Linear Equations with constant coefficients and second member *not* zero.

There are two methods of solution :

FIRST. METHOD OF VARIABLE PARAMETERS. — Solve the equation by Art. 43, regarding the second member as zero.

Supposing it to be of the n th order, this value of y will contain n arbitrary constants. Derive from it the successive differential coefficients, $\frac{dy}{dx}$, $\frac{d^2 y}{dx^2}$, ... $\frac{d^{n-1} y}{dx^{n-1}}$; then differentiate the values of y , $\frac{dy}{dx}$, $\frac{d^2 y}{dx^2}$, ... $\frac{d^{n-1} y}{dx^{n-1}}$, regarding the arbitrary constants alone as variable, and place these n results equal to zero, except the last, which put equal to the second member of the given equation. These n conditions will determine expressions for the n arbitrary constants, which are to be substituted in the original expression for y .

SECOND METHOD. — By successively differentiating the given equation, obtain, either directly or by elimination, a new differential equation of a higher order with the second member zero. Solve this by Art. 43, and determine the values of the superfluous constants so as to satisfy the given differential equation. In this last work of determining the superfluous constants all the other constants may be regarded as zero.

48. Solve the following equations :

1. $\frac{d^2 y}{dx^2} - 7 \frac{dy}{dx} + 12y = x.$
2. $\frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = a.$
3. $\frac{d^2 y}{dx^2} - a^2 y = x + 1.$
4. $\frac{d^3 y}{dx^3} - 2 \frac{d^2 y}{dx^2} + \frac{dy}{dx} = e^x.$
5. $\frac{d^4 y}{dx^4} - a^4 y = x^3.$
6. $\frac{d^2 y}{dx^2} + n^2 y = 1 + x + x^2.$
7. $\frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + a^2 y = e^x$; also when $a = 1.$
8. $\frac{d^2 y}{dx^2} + n^2 y = \cos ax$; also when $a = n.$
9. $\frac{d^2 y}{dx^2} - 5 \frac{dy}{dx} + 6y = e^{nx}$; also when $n = 2$, or $n = 3.$
10. $\frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y = xe^{nx}.$
11. $\frac{d^2 y}{dx^2} - 9 \frac{dy}{dx} + 20y = x^2 e^{3x}.$
12. $\frac{d^2 y}{dx^2} + 4y = x \sin^2 x.$

$$13. \quad \frac{d^4 y}{dx^4} + 2 \frac{d^2 y}{dx^2} + y = x^2 \cos ax; \quad \text{also when } a = 1.$$

$$14. \quad \frac{d^3 y}{dx^3} - 2 \frac{dy}{dx} + 4y = e^x \cos x.$$

49. Linear equations of the form

$$(a+bx)^n \frac{d^n y}{dx^n} + A_1(a+bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + A_{n-1}(a+bx) \frac{dy}{dx} + A_n y = X,$$

where A_1, A_2, \dots, A_n are constants, and X a function of x alone.

Put $a+bx = e^t$, and change the independent variable from x to t . The new differential equation between y and t will be linear with *constant* coefficients, and may be solved by Art. 47.

50. Solve the following equations :

$$1. \quad x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 3y.$$

$$2. \quad (x+a)^2 \frac{d^2 y}{dx^2} - 4(x+a) \frac{dy}{dx} + 6y = x.$$

$$3. \quad (a+bx)^2 \frac{d^2 y}{dx^2} + b(a+bx) \frac{dy}{dx} + b^2 y = 0.$$

$$4. \quad x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + 2y = x \log x.$$

$$5. \quad (2x-1)^3 \frac{d^3 y}{dx^3} + (2x-1) \frac{dy}{dx} - 2y = 0.$$

$$6. \quad 16(x+1)^4 \frac{d^4 y}{dx^4} + 96(x+1)^3 \frac{d^3 y}{dx^3} + 104(x+1)^2 \frac{d^2 y}{dx^2} \\ + 8(x+1) \frac{dy}{dx} + y = x^2 + 4x + 3.$$

$$7. \quad x^4 \frac{d^4 y}{dx^4} + 6x^3 \frac{d^3 y}{dx^3} + 9x^2 \frac{d^2 y}{dx^2} + 3x \frac{dy}{dx} + y = (1 + \log x)^2.$$

$$8. \quad x^2 \frac{d^2 y}{dx^2} - (2m-1)x \frac{dy}{dx} + (m^2 + n^2)y = n^2 x^m \log x.$$

CHAPTER VIII.

SOME SPECIAL FORMS OF DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

51. Form, $\frac{d^n y}{dx^n} = X$, where X is a function of x alone.

The expression for y is found by integrating X successively n times with regard to x . Or solve by Art. 47.

52. Solve the following equations :

1. $x \frac{d^3 y}{dx^3} = 2.$

2. $\frac{d^4 y}{dx^4} = \frac{1}{(x+a)^2}.$

3. $\frac{d^n y}{dx^n} = x^n.$

4. $\frac{d^4 y}{dx^4} = x \cos x.$

5. $e^x \frac{d^4 y}{dx^4} + 4 \cos x = 0.$

6. $\frac{d^n y}{dx^n} = x e^x.$

7. $\frac{d^3 y}{dx^3} = \sin^3 x.$

53. Form, $\frac{d^2 y}{dx^2} = Y$, where Y is a function of y only.

Multiplying both members by $2 \frac{dy}{dx}$, and integrating, we have
 $\left(\frac{dy}{dx}\right)^2 = 2 \int Y dy + c_1.$ Therefore $x = \int \frac{dy}{(2 \int Y dy + c_1)^{\frac{1}{2}}} + c_2.$

54. Solve the following equations :

$$1. \quad \frac{d^2 y}{dx^2} = a^2 y.$$

$$2. \quad \frac{d^2 y}{dx^2} = -a^2 y.$$

$$3. \quad y^3 \frac{d^2 y}{dx^2} = a.$$

$$4. \quad \frac{d^2 y}{dx^2} = e^{ny}.$$

$$5. \quad \frac{d^2 y}{dx^2} = \frac{1}{\sqrt{ay}}.$$

55. *Equations not containing y directly.*

By assuming the differential coefficient of the lowest order in the given equation equal to z , and consequently the other differential coefficients equal to the successive differential coefficients of z with respect to x , we shall obtain a new differential equation between z and x of a lower order than the given equation.

56. Solve the following equations :

$$1. \quad x \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 0.$$

$$2. \quad \frac{d^2 y}{dx^2} = a^2 + b^2 \left(\frac{dy}{dx} \right)^2.$$

$$3. \quad \frac{dy}{dx} - x \frac{d^2 y}{dx^2} = f \left(\frac{d^2 y}{dx^2} \right).$$

$$4. \quad a^2 \left(\frac{d^2 y}{dx^2} \right)^2 = 1 + \left(\frac{dy}{dx} \right)^2.$$

$$5. \quad a^2 \left(\frac{d^2 y}{dx^2} \right)^2 = \left[1 + \left(\frac{dy}{dx} \right)^2 \right]^3.$$

$$6. \quad (1 + x^2) \frac{d^2 y}{dx^2} + 1 + \left(\frac{dy}{dx} \right)^2 = 0.$$

7. $(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 2.$
8. $2x \frac{d^3 y}{dx^3} \cdot \frac{d^2 y}{dx^2} = \left(\frac{d^2 y}{dx^2} \right)^2 - \alpha^2.$
9. $\frac{d^3 y}{dx^3} \cdot \frac{d^2 y}{dx^2} = \left(1 - \frac{d^3 y}{dx^3} \right) \left[1 + \left(\frac{d^2 y}{dx^2} \right)^2 \right]^{\frac{1}{2}}.$
10. $\frac{d^3 y}{dx^3} \left(\frac{dy}{dx} \right)^3 = 1.$

57. *Equations not containing x directly.*

By assuming $\frac{dy}{dx} = z$, and consequently

$$\frac{d^2 y}{dx^2} = z \frac{dz}{dy}, \quad \frac{d^3 y}{dx^3} = z^2 \frac{d^2 z}{dy^2} + z \left(\frac{dz}{dy} \right)^2, \text{ etc.,}$$

changing the independent variable from x to y , we shall obtain a new differential equation between z and y of a lower order than the given equation.

58. Solve the following equations :

1. $y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx} \right)^2 = 1.$
2. $y(1 - \log y) \frac{d^2 y}{dx^2} + (1 + \log y) \left(\frac{dy}{dx} \right)^2 = 0.$
3. $y \frac{d^2 y}{dx^2} + \left[\left(\frac{dy}{dx} \right)^2 + \alpha^2 \left(\frac{d^2 y}{dx^2} \right)^2 \right]^{\frac{1}{2}} = \left(\frac{dy}{dx} \right)^2.$
4. $\left(\frac{dy}{dx} \right)^2 - y \frac{d^2 y}{dx^2} = \frac{dy}{dx} \cdot f \left[\left(\frac{dy}{dx} \right)^{-1} \cdot \frac{d^2 y}{dx^2} \right].$
5. $y \frac{d^2 y}{dx^2} - \left(\frac{dy}{dx} \right)^2 = y^2 \log y.$
6. $y^2 \left(\frac{d^2 y}{dx^2} \right)^2 + n \left(\frac{dy}{dx} \right)^4 = y^2 \frac{dy}{dx} \cdot \frac{d^3 y}{dx^3} + n y \left(\frac{dy}{dx} \right)^2 \frac{d^2 y}{dx^2}.$

CHAPTER IX.

SIMULTANEOUS DIFFERENTIAL EQUATIONS.

59. *Simultaneous differential equations of the first order.*

There should be n given equations between $n + 1$ variables. Selecting one of these for the independent variable, we may, by differentiating the given equations a sufficient number of times, eliminate all but one of the dependent variables and their differential coefficients. The resulting differential equation between two variables must be solved by the methods previously given, and from its primitive and the given equations may be obtained the values of the other dependent variables. The complete solution will consist of n equations containing n arbitrary constants.

In general, if we differentiate the given equations $n - 1$ times successively, we shall have in all n^2 equations, which are just sufficient for the elimination of $n - 1$ variables, together with their $n(n - 1)$ differential coefficients. Shorter processes for the elimination will frequently suggest themselves in special cases.

60. Solve the following simultaneous equations :

$$1. \quad \begin{cases} \frac{dx}{dt} + 4x + \frac{y}{4} = 0, \\ \frac{dy}{dt} + 3y - x = 0. \end{cases}$$

$$2. \quad -dx = \frac{dy}{3y + 4z} = \frac{dz}{2y + 5z}.$$

$$3. \quad \frac{dx}{y - 7x} = \frac{-dy}{2x + 5y} = dt.$$

$$4. \quad \begin{cases} a \frac{dz}{dx} + n^2 y = e^z, \\ \frac{dy}{dx} + az = 0. \end{cases}$$

$$5. \quad \begin{cases} \frac{dx}{dt} + 5x + y = e^t, \\ \frac{dy}{dt} + 3y - x = e^{2t}. \end{cases}$$

$$6. \quad \frac{dx}{2y - 5x + e^t} = \frac{dy}{x - 6y + e^{2t}} = dt.$$

$$7. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 44x + 49y = t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + 34x + 38y = e^t. \end{cases}$$

$$8. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 11x + 31y = e^t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + 8x + 24y = e^{2t}. \end{cases}$$

$$9. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 2x + 31y = e^t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + x + 24y = 3. \end{cases}$$

$$10. \quad \begin{cases} \frac{dx}{dt} + \frac{2}{t}(x - y) = 1, \\ \frac{dy}{dt} + \frac{1}{t}(x + 5y) = t. \end{cases}$$

$$11. \quad \begin{cases} t dx = (t - 2x) dt, \\ t dy = (tx + ty + 2x - t) dt. \end{cases}$$

$$12. \quad \begin{cases} \frac{dx}{dt} = ny - mz, \\ \frac{dy}{dt} = lz - nx, \\ \frac{dz}{dt} = mx - ly. \end{cases}$$

$$13. \quad \begin{cases} lt \frac{dx}{dt} = mn (y - z), \\ mt \frac{dy}{dt} = nl (z - x), \\ nt \frac{dz}{dt} = lm (x - y). \end{cases}$$

61. *Simultaneous differential equations of an order higher than the first.*

By differentiating the given equations a sufficient number of times, we may eliminate all but one of the dependent variables and their differential coefficients, and thus obtain a differential equation between two variables, which must be solved by the appropriate methods. Its primitive, together with the given equation, will enable us to determine the values of the other dependent variables. The general solution will contain a number of arbitrary constants equal to the sum of the highest orders of differential coefficients in the several given equations.

62. Solve the following simultaneous equations :

$$1. \quad \begin{cases} \frac{d^2x}{dt^2} + n^2x = 0, \\ \frac{d^2y}{dt^2} - n^2x = 0. \end{cases}$$

$$2. \quad \begin{cases} \frac{d^2x}{dt^2} - 3x - 4y + 3 = 0, \\ \frac{d^2y}{dt^2} + x + y + 5 = 0. \end{cases}$$

$$3. \quad \begin{cases} \frac{d^2 x}{dt^2} - 3x - 4y + 3 = 0, \\ \frac{d^2 y}{dt^2} + x - 8y + 5 = 0. \end{cases}$$

$$4. \quad \begin{cases} \frac{d^2 x}{dt^2} + n^2 y = 0, \\ \frac{d^2 y}{dt^2} - n^2 x = 0. \end{cases}$$

$$5. \quad \begin{cases} 2 \frac{d^2 y}{dx^2} - \frac{dz}{dx} - 4y = 2x, \\ 2 \frac{dy}{dx} + 4 \frac{dz}{dx} - 3z = 0. \end{cases}$$

$$6. \quad \begin{cases} \frac{d^3 y}{dx^3} + 4 \frac{d^2 z}{dx^2} + (5 - n^2) \frac{dy}{dx} + 2(1 - n^2)z = 0, \\ \frac{d^3 z}{dx^3} + 4 \frac{d^2 y}{dx^2} + (5 - n^2) \frac{dz}{dx} + 2(1 - n^2)y = 0. \end{cases}$$

$$7. \quad \begin{cases} \frac{d^4 y}{dx^4} - 4 \frac{d^3 z}{dx^3} + 4 \frac{d^2 y}{dx^2} - y = 0, \\ \frac{d^4 z}{dx^4} - 4 \frac{d^3 y}{dx^3} + 4 \frac{d^2 z}{dx^2} - z = 0. \end{cases}$$

CHAPTER X.

GEOMETRICAL EXAMPLES.

63. Expressions involved in the examples, p representing $\frac{dy}{dx}$, and q representing $\frac{d^2y}{dx^2}$.

$$\text{Subtangent} = \frac{y}{p}. \quad \text{Subnormal} = py.$$

$$\text{Normal} = y \sqrt{1 + p^2}. \quad \frac{ds}{dx} = \sqrt{1 + p^2}.$$

$$\text{Intercept of tangent on axis of } X = x - \frac{y}{p}.$$

$$\text{Intercept of tangent on axis of } Y = y - px.$$

$$\text{Radius of curvature} = \mp \frac{(1 + p^2)^{\frac{3}{2}}}{q}.$$

1. Find the curve whose subtangent varies as (is n times) the abscissa.
2. Find the curve whose subnormal is constant and equal to $2a$.
3. Find the curve whose normal is equal to the square of the ordinate.
4. Find the curve for which $s = mx^2$.
5. Find the curve for which $s^2 = y^2 - a^2$.

The *orthogonal trajectory* of a series of curves is a curve that intersects them all at right angles.

Describe the curves represented by the following equations, and find their orthogonal trajectories:

6. $y = mx$, m being the variable parameter.

7. $y^2 = 2ax - x^2$, a being the variable parameter.
8. $y^2 = 4ax$, a being the variable parameter.
9. $xy = k^2$, k being the variable parameter.
10. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, b being the variable parameter.
11. $x^2 + m^2y^2 = m^2a^2$, a being the variable parameter.
12. $\frac{x^2}{b^2 + h^2} + \frac{y^2}{b^2} = 1$, b being the variable parameter.

The three following examples require the singular solution :

13. Find the curve such that the sum of the intercepts of the tangent on the axes of X and Y is constant and equal to a .
14. Find the curve such that the part of the tangent between the axes of X and Y is constant and equal to a .
15. Find the curve such that the area of the right triangle formed by the tangent with the axes of X and Y is constant and equal to a^2 .

The following examples require the solution of differential equations of the second order :

16. Find the curve such that the length of the arc measured from some fixed point of it is equal to the intercept of the tangent on the axis of X .
17. Find the curve whose radius of curvature varies as (is n times) the cube of the normal.
18. Find the curve whose radius of curvature is equal to the normal ; first, when the two have the same direction ; second, when they have opposite directions.
19. Find the curve whose radius of curvature is equal to twice the normal ; first, when the two have the same direction ; second, when they have opposite directions.

ANSWERS.



Art. 3. $\left(p = \frac{dy}{dx}\right)$

1. $y(1+x) + px(1-y) = 0$.
2. $(x^2+1)pxy = y^2+1$.
3. $\tan x = p \tan y$.
4. $(1+x^2)p + y = \tan^{-1}x$.
5. $(y \log x - 1)y = px$.
6. $y = px + p - p^3$.
7. $xp^2 = a$.
8. $p^2 + 2py \cot x = y^2$.
9. $x^2p^2 = 1 + p^2$.

Art. 5.

1. $\frac{d^2y}{dx^2} + a^2y = 0$.
2. $\frac{d^2y}{dx^2} - a^2y = 0$.
3. $\frac{d^2y}{dx^2} - 2a \frac{dy}{dx} + a^2y = 0$.
4. $x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 3y$.
5. $\frac{d^2y}{dx^2} + n^2y = \cos ax$.

Art. 7.

1. $\frac{d^3y}{dx^3} = 7 \frac{dy}{dx} - 6y$.
2. $\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} + \frac{dy}{dx} = 3y$.
3. $\frac{d^3y}{dx^3} - 2 \frac{d^2y}{dx^2} + \frac{dy}{dx} = e^x$.
4. $\frac{d^4y}{dx^4} - 3 \frac{d^3y}{dx^3} + 3 \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$.
5. $\frac{d^4y}{dx^4} - a^4y = x^3$.

Art. 9.

1. $\log(xy) + x - y = c$.
2. $\frac{x+y}{xy} + \log \frac{y}{x} = c$.
3. $(1+x^2)(1+y^2) = cx^2$.
4. $x^2y = ce^{\frac{y}{x}}$.

$$5. \log [(y + \sqrt{1+y^2}) \sqrt{1+y^2}] = \frac{x}{\sqrt{1+x^2}} + c.$$

$$6. \cos y = c \cos x.$$

$$8. \sin^2 x + \sin^2 y = c.$$

$$7. \tan x \tan y = c.$$

$$9. xy - 1 = c(x + y + 1).$$

Art. 11.

$$1. y = ce^{-\frac{x}{y}}.$$

$$5. x = ce^{-\sin \frac{y}{x}}.$$

$$2. y = ce^{-\sqrt{\frac{x}{y}}}.$$

$$6. (y+x)^2 (y+2x)^3 = c.$$

$$3. y = ce^{\frac{y}{x}}.$$

$$7. \log (x^2 + y^2) = 2 \tan^{-1} \frac{y}{x} + c.$$

$$4. x^2 = c^2 + 2cy.$$

$$8. xy \cos \frac{y}{x} = c.$$

$$9. (1), \log (x^2 - mxy + y^2) + \frac{2m}{\sqrt{4-m^2}} \tan^{-1} \frac{2y-mx}{x\sqrt{4-m^2}} = c.$$

$$(2), x - y = ce^{\frac{x}{y-2}}.$$

$$(3), \frac{(2y - mx + x\sqrt{m^2 - 4})^{m - \sqrt{m^2 - 4}}}{(2y - mx - x\sqrt{m^2 - 4})^{m + \sqrt{m^2 - 4}}} = c.$$

$$10. y \sin a - x \cos a + \sqrt{x^2 + y^2} = c(x^2 + y^2).$$

Art. 13.

$$1. (y - x + 1)^2 (y + x - 1)^5 = c.$$

$$2. x + 2y + \log (2x + y - 1) = c.$$

$$3. x + 5y + 2 = c(x - y + 2)^4.$$

$$4. 4x - 8y = \log (4x + 8y + 5) + c.$$

$$5. \log [2(3x-1)^2 + (3y-5)^2] - \sqrt{2} \tan^{-1} \frac{\sqrt{2}(3x-1)}{3y-5} = c.$$

Art. 15.

1. $y = cx^a + \frac{x}{1-a} - \frac{1}{a}$.
2. $y = ax + cx\sqrt{1-x^2}$.
3. $y = \frac{x}{\sqrt{1-x^2}} + ce^{\frac{x}{\sqrt{1-x^2}}}$.
4. $y = \sin x - 1 + ce^{-\sin x}$.
5. $x = \tan^{-1}y - 1 + ce^{-\tan^{-1}y}$.
6. $(x + \sqrt{a^2 + x^2})y = a^2 \log(x + \sqrt{a^2 + x^2}) + c$.
7. $y\sqrt{1+x^2} = \log \frac{\sqrt{1+x^2} - 1}{x} + c$.
8. $y = ce^{-\phi} + \phi - 1$.

Art. 17.

1. $y = (c\sqrt{1-x^2} - a)^{-1}$.
2. $y^3 = ce^{ax} - \frac{x+1}{a} - \frac{1}{a^2}$.
3. $y = \left[ce^{2x^2} + \frac{a}{2}(2x^2 + 1) \right]^{-\frac{1}{2}}$.
4. $x = \frac{e^{\frac{y^2}{2}}}{(2 - y^2)e^{\frac{y^2}{2}} + c}$.
5. $y^{-n+1} = ce^{(n-1)\sin x} + 2\sin x + \frac{2}{n-1}$.
6. $y = (cx + \log x + 1)^{-1}$.
7. $x = \frac{(n+2)y^2}{ay^{n+2} + c}$.
8. $y = \frac{\tan x + \sec x}{\sin x + c}$.
9. $(4b^2 + 1)y^2 = 2a(\sin x + 2b\cos x) + ce^{-2bx}$.

Art. 19.

1. $x^4 + 6x^2y^2 + y^4 = c$.
2. $x^3 - 6x^2y - 6xy^2 + y^3 = c$.
3. $x^2 - y^2 = cx$.
4. $x^2 - y^2 = cy^3$.
5. $x^2 + y^2 + 2\tan^{-1}\frac{y}{x} = c$.
6. $y^2 = c^2 - 2cx$.
7. $x^2 + y^2 + 2\sin^{-1}\frac{x}{y} = c$.
8. $x + ye^{\frac{x}{y}} = c$.

$$9. e^x(x^2 + y^2) = c.$$

$$10. \cos(mx + ny) + \sin(nx + my) = c.$$

$$11. \sqrt{1 + x^2 + y^2} + \tan^{-1} \frac{x}{y} = c.$$

$$12. (1), \log \frac{x^a \sqrt{g} + y \sqrt{b}}{x^a \sqrt{g} - y \sqrt{b}} = \frac{2x^a \sqrt{bg}}{a} + c.$$

$$(2), \tan^{-1} \frac{y \sqrt{b}}{x^a \sqrt{k}} + \frac{x^a \sqrt{bk}}{a} = c.$$

$$(3), x^a \left(\frac{1}{a} - \frac{1}{by} \right) = c.$$

$$(4), \frac{\sqrt{g} + y \sqrt{b}}{\sqrt{g} - y \sqrt{b}} = cx^{2\sqrt{bg}}.$$

$$(5), \frac{x^a}{a} - \frac{y}{gx^a} = c.$$

Art. 22.

$$1. x^2 + y^2 = c(x + y).$$

$$4. x^2 + 2y^2 = c\sqrt{x^2 + y^2}.$$

$$2. x^2 - y^2 + xy = c.$$

$$5. y = cx.$$

$$3. y = cx.$$

Art. 24.

$$1. x = cy e^{\frac{1}{xy}}.$$

$$4. \frac{2}{\sqrt{xy}} = \log \frac{cx}{y}.$$

$$2. y = ce^{xy}.$$

$$5. xy = c.$$

$$3. xy - \frac{1}{xy} = \log cy^2.$$

$$6. xye^{xy} = \log \frac{cy}{x}.$$

$$7. xy + \log \sin(xy) = \log \frac{cx}{y}.$$

Art. 26.

$$1. e^x(x^2 + y^2) = c.$$

$$3. x^2 - y^2 = cx.$$

$$2. x^2 - y^2 = cy^3.$$

$$4. y\sqrt{1-x^2} + x(1-y) = c.$$

$$5. \sin x \cos y + y \tan 2x = c. \quad 6. \sin^2 x \sin 2(x - y) = c.$$

$$7. y = e^{-\int P dx} \left(\int Q e^{\int P dx} dx + c \right).$$

Art. 28.

$$1. (x - a)(y - b)(z - c) = c'. \quad 7. z = ce^{\frac{x}{y}}.$$

$$2. x^2 + 2y^2 - 6xy - 2xz + z^2 = c. \quad 8. xy = c(a + z).$$

$$3. yz + zx + xy = c. \quad 9. x = \frac{z}{y + a} + c.$$

$$4. xyz = c. \quad 10. y(x + z) = c(y + z).$$

$$5. e^x(y + z) = c. \quad 11. e^{x^2}(x + y + z^2) = c.$$

$$6. \frac{a}{x} + \frac{b}{y} + \frac{c}{z} = c'. \quad 12. \frac{y + z}{x} + \frac{z + x}{y} = c.$$

Art. 30.

$$1. (y - 2x + c)(y - 3x + c) = 0, \text{ or } (5x - 2y + c)^2 = x^2,$$

$$2. (y + c)^2 = a^2(\log x)^2. \quad 3. (y + c)^2 = 4ax.$$

$$4. (y + c)^2 = (\sqrt{x - x^2} + \sin^{-1} \sqrt{x})^2.$$

$$5. (xy + c)(x^2y + c) = 0.$$

$$6. (x^2 - 2y + c)[e^x(x + y - 1) + c] = 0.$$

$$7. (y + c)(y + x^2 + c)(xy + cy + 1) = 0.$$

$$8. (x^3 - 3y + c)(e^{\frac{x^2}{2}} + cy)(xy + cy + 1) = 0.$$

$$9. y^2 \sin^2 x + 2cy + c^2 = 0.$$

Art. 32.

$$1. \text{ Eliminate } p \text{ by means of } x = \frac{p}{\sqrt{1 - p^2}} (c + a \sin^{-1} p).$$

$$2. \text{ Eliminate } p \text{ by means of } x(p - 1)^2 = \log p^2 - 2p + c.$$

$$3. \text{ Eliminate } p \text{ by means of } x = \frac{p}{\sqrt{1 + p^2}} \left(c + \frac{a}{p} + a \tan^{-1} p \right).$$

$$4. y = cx + c - c^2.$$

$$5. x = a \log (ay \pm \sqrt{a^2 + y^2 - 1}) + \log (y \mp \sqrt{a^2 + y^2 - 1}) + c.$$

$$6. x \pm \sqrt{a^2 + 4by} = a \log (a \pm \sqrt{a^2 + 4by}) + c.$$

$$7. c(2y \pm x\sqrt{x^2 + y})^{\sqrt{x}} = \frac{\pm \sqrt{x^2 + y} - x(\sqrt{17} - 1)}{\pm \sqrt{x^2 + y} + x(\sqrt{17} + 1)}.$$

$$8. \left[y\sqrt{y^2 - x^2} - x^2 \log \frac{y + \sqrt{y^2 - x^2}}{x} \right]^2 = (y^2 + x^2 \log cx^2)^2.$$

$$9. 4(x+c)^3 + (x+c)^2 - 18y(x+c) - 27y^2 - 4y = 0.$$

Art. 34.

$$1. y^2 = 2cx + c^2.$$

$$2. x + 1 = \pm \sqrt{2y + c} + \log (\pm \sqrt{2y + c} - 1).$$

$$3. (y + c)^2 = \frac{(x + a)^2}{x^2 + 2ax}.$$

$$4. e^{2y} + 2cxe^y + c^2 = 0.$$

$$5. 2y + c = \frac{ax^2 \pm x\sqrt{a^2 + x^2 - 1}}{a^2 - 1} + \log (x \pm \sqrt{a^2 + x^2 - 1});$$

$$\text{when } a=1, \quad 4y = x^2 - \log (cx^2).$$

$$6. b^2(6y + c)^2 + (6abx + a^3)(6y + c) - 3a^2x^2 - 16bx^3 = 0.$$

$$7. c(nx^2 + 2y^2 \pm x\sqrt{n^2x^2 + 4my^2})^n = [(2m - n)x \pm \sqrt{n^2x^2 + 4my^2}]^{2m}.$$

Art. 36.

$$1. (x^2 - y^2 + c)(x^2 - y^2 + cx^4) = 0.$$

$$2. (\sqrt{x} - \sqrt{y} + c)(\sqrt{x} - \sqrt{y} + cx) = 0.$$

$$3. x^4 + 2cx\sqrt{3y^2 - x^2} - c^2 = 0.$$

$$4. (y - x)^{1 \pm \sqrt{2}} = c(\sqrt{y} + \sqrt{x})^2.$$

$$5. x^{2k} - 2cx^{k-1}\sqrt{y^2 + nx^2} + nc^2 = 0, \quad \text{where } k = \sqrt{\frac{n-1}{n}}.$$

Art. 38.

1. $y = cx + \frac{m}{c}$.
2. $y = cx + c - c^3$.
3. $(y - cx)^2 = 1 + c^2$.
4. $y^2 = cx + \frac{c^3}{8}$.
5. $4y^2 = 2c(2x - b) + ac^2$.
6. $y^2 = cx^2 + c^2$.
7. $e^y = ce^x + c^3$.
8. $y^2 - cx^2 = -\frac{ch^2}{c+1}$.

Art. 42.

Complete Primitives and Singular Solutions :

1. $y = cx + \frac{m}{c}, \quad y^2 = 4mx.$
2. $(y - cx)^2 = 1 - c^2, \quad y^2 - x^2 = 1.$
3. $y = c(x - c)^2, \quad y = \frac{4x^3}{27}.$
4. $y = c(x - 1) - c^2, \quad 4y = (x - 1)^2.$
5. $y^2 - 2cx + c^2 = 0, \quad y^2 = x^2.$
6. $(y - cx)^2 = 4c, \quad xy = 1.$
7. $(y - cx)(mc - n) = mnc, \quad \left(\frac{x}{m}\right)^{\frac{1}{2}} \pm \left(\frac{y}{n}\right)^{\frac{1}{2}} = 1.$

Art. 44.

1. $y = c_1 e^{ax} + c_2 e^{-ax}.$
2. $y = c_1 e^{3x} + c_2 e^{4x}.$
3. $y = c_1 e^{\frac{x}{a}} + c_2.$
4. $y = c_1 e^{3x} + c_2 e^{\frac{x}{3}}.$
5. $ye^{2x} = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}}.$
6. $y = c_1 e^{\frac{ax}{b}} + c_2 e^{\frac{bx}{a}}.$
7. $y = c_1 e^{2x} + c_2 e^{-2x} + c_3.$
8. $y = c_1 e^{3x} + c_2 e^{-2x} + c_3.$
9. $y = c_1 e^{2x} + c_2 e^{-3x} + c_3 e^x.$
10. $y = c_1 e^{3x} + c_2 e^{-3x} + c_3 e^{x\sqrt{3}} + c_4 e^{-x\sqrt{3}}.$
11. $y = c_1 e^{(a-b)x} + c_2 e^{(b-a)x} + c_3 e^{(a+b)x} + c_4 e^{-(a+b)x} + c_5.$

Art. 45.

1. $y = c_1 \sin x + c_2 \cos x$.
2. $y = e^{3x}(c_1 \sin 2x + c_2 \cos 2x)$.
3. When $a > b$, $y = e^{ax}(c_1 e^{\sqrt{a^2-b^2}x} + c_2 e^{-\sqrt{a^2-b^2}x})$;
when $a < b$, $y = e^{ax}(c_1 \sin x \sqrt{b^2-a^2} + c_2 \cos x \sqrt{b^2-a^2})$.
4. $y = e^{2abx}[c_1 \sin(a^2-b^2)x + c_2 \cos(a^2-b^2)x]$.
5. $y = a^x(c_1 \sin x + c_2 \cos x)$.
6. $y = c_1 \sin x \sqrt{2} + c_2 \cos x \sqrt{2} + c_3$.
7. $y = c_1 e^x + e^{-\frac{x}{2}}\left(c_2 \sin \frac{x\sqrt{3}}{2} + c_3 \cos \frac{x\sqrt{3}}{2}\right)$.
8. $ye^x = c_1 e^{2x} + c_2 \sin x \sqrt{2} + c_3 \cos x \sqrt{2}$.
9. $y = c_1 e^x + c_2 e^{-x} + c_3 \sin x + c_4 \cos x$.
10. $y = c_1 e^{xv^2} + c_2 e^{-xv^2} + c_3 \sin 2x + c_4 \cos 2x$.
11. $y = e^{ax}(c_1 \sin ax + c_2 \cos ax) + e^{-ax}(c_3 \sin ax + c_4 \cos ax)$.
12. $y = c_1 e^x + c_2 e^{-x} + (c_3 e^{\frac{x}{2}} + c_4 e^{-\frac{x}{2}}) \sin \frac{x\sqrt{3}}{2}$
 $+ (c_5 e^{\frac{x}{2}} + c_6 e^{-\frac{x}{2}}) \cos \frac{x\sqrt{3}}{2}$.
13. $y = c_1 \sin x + c_2 \cos x + e^{\frac{x\sqrt{3}}{2}}\left(c_3 \sin \frac{x}{2} + c_4 \cos \frac{x}{2}\right)$
 $+ e^{-\frac{x\sqrt{3}}{2}}\left(c_5 \sin \frac{x}{2} + c_6 \cos \frac{x}{2}\right)$.
14. $y = c_1 e^x + c_2 e^{-x} + c_3 \sin x + c_4 \cos x + e^{\frac{x}{\sqrt{2}}}\left(c_5 \sin \frac{x}{\sqrt{2}} + c_6 \cos \frac{x}{\sqrt{2}}\right)$
 $+ e^{-\frac{x}{\sqrt{2}}}\left(c_7 \sin \frac{x}{\sqrt{2}} + c_8 \cos \frac{x}{\sqrt{2}}\right)$.

Art. 46.

1. $y = (c_1 + c_2 x) e^{ax}$.
2. $y = c_1 + c_2 x$.
3. $y = c_1 e^{4x} + c_2 + c_3 x$.
4. $y = c_1 e^{-x} + (c_2 + c_3 x) e^{2x}$.

5. $y = c_1 e^{-x} + (c_2 + c_3 x) e^x.$
 6. $y = (c_1 + c_2 x) \cos nx + (c_3 + c_4 x) \sin nx.$
 7. $y = (c_1 + c_2 x + c_3 x^2) e^x + c_4.$
 8. $y = e^x [(c_1 + c_2 x) \sin 2x + (c_3 + c_4 x) \cos 2x].$
 9. $y = c_1 + c_2 x + c_3 x^2 \dots + c_{n-2} x^{n-3} + c_{n-1} \sin x + c_n \cos x.$

Art. 48.

1. $y = c_1 e^{3x} + c_2 e^{4x} + \frac{12x + 7}{141}.$
 2. $y = c_1 \sin x + c_2 \cos x + (c_3 + c_4 x) e^x + a.$
 3. $y = c_1 e^{ax} + c_2 e^{-ax} - \frac{x + 1}{a^2}.$
 4. $y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x + c_3.$
 5. $y = c_1 e^{ax} + c_2 e^{-ax} + c_3 \sin ax + c_4 \cos ax - \frac{x^3}{a^4}.$
 6. $y = c_1 \sin nx + c_2 \cos nx + \frac{1 + x + x^2}{n^2} - \frac{2}{n^4}.$
 7. $y = (c_1 + c_2 x) e^{ax} + \frac{e^x}{(a-1)^2};$
 when $a = 1, \quad y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x.$
 8. $y = c_1 \sin nx + c_2 \cos nx + \frac{\cos ax}{n^2 - a^2};$
 when $a = n, \quad y = c_1 \sin nx + c_2 \cos nx + \frac{x \sin nx}{2n}.$
 9. $y = c_1 e^{2x} + c_2 e^{3x} + \frac{e^{nx}}{n^2 - 5n + 6};$
 when $n = 2, \quad y = (c_1 - x) e^{2x} + c_2 e^{3x};$
 when $n = 3, \quad y = c_1 e^{2x} + (c_2 + x) e^{3x}.$
 10. $y = c_1 e^x + c_2 e^{2x} + \frac{x e^{nx}}{n^2 - 3n + 2} - \frac{(2n - 3) e^{nx}}{(n^2 - 3n + 2)^2}.$

$$11. y = c_1 e^{4x} + c_2 e^{5x} + \frac{2x^2 + 6x + 7}{4} e^{3x}.$$

$$12. y = \left(c_1 - \frac{x^2}{16}\right) \sin 2x + \left(c_2 - \frac{x}{32}\right) \cos 2x + \frac{x}{8}.$$

$$13. y = (c_1 + c_2 x) \sin x + (c_3 + c_4 x) \cos x \\ + \left[\frac{x^2}{(a^2 - 1)^2} - \frac{4(5a^2 + 1)}{(a^2 - 1)^4} \right] \cos ax - \frac{8ax}{(a^2 - 1)^3} \sin ax;$$

$$\text{when } a = 1, \quad y = \left(c_1 + c_2 x + \frac{x^3}{12} - \frac{x}{4}\right) \sin x \\ + \left(c_3 + c_4 x - \frac{x^4}{48} + \frac{3x^2}{16} + \frac{3}{8}\right) \cos x.$$

$$14. y = c_1 e^{-2x} + \left(c_2 + \frac{33}{400} - \frac{x}{20}\right) e^x \cos x + \left(c_3 - \frac{3}{80} + \frac{3x}{20}\right) e^x \sin x.$$

Art. 50.

$$1. y = c_1 x^3 + \frac{c_2}{x}.$$

$$2. y = c_1(x+a)^2 + c_2(x+a)^3 + \frac{3x+2a}{6}.$$

$$3. y = c_1 \sin \log(a+bx) + c_2 \cos \log(a+bx).$$

$$4. y = x(c_1 \sin \log x + c_2 \cos \log x + \log x).$$

$$5. y = (2x-1) \left[c_1 + c_2 (2x-1)^{\frac{\sqrt{3}}{2}} + c_3 (2x-1)^{-\frac{\sqrt{3}}{2}} \right].$$

$$6. y = [c_1 + c_2 \log(x+1)] \sqrt{x+1} + \frac{c_3 + c_4 \log(x+1)}{\sqrt{x+1}}.$$

$$7. y = (c_1 + c_2 \log x) \sin \log x + (c_3 + c_4 \log x) \cos \log x + (\log x)^2 \\ + 2 \log x - 3.$$

$$8. y = x^m (c_1 \sin \log x^n + c_2 \cos \log x^n + \log x).$$

Art. 52.

$$1. y = c_1 + c_2 x + c_3 x^2 + x^2 \log x.$$

$$2. y = c_1 + c_2 x + c_3 x^2 + c_4 x^3 - (x+a)^2 \log \sqrt{x+a}.$$

3. $y = c_1 + c_2 x + c_3 x^2 \dots + c_n x^{n-1} + \frac{m x^{m+n}}{m+n}.$
4. $y = c_1 + c_2 x + c_3 x^2 + c_4 x^3 + x \cos x - 4 \sin x.$
5. $y = c_1 + c_2 x + c_3 x^2 + c_4 x^3 + e^{-x} \cos x.$
6. $y = c_1 + c_2 x + c_3 x^2 \dots + c_n x^{n-1} + (x-n)e^x.$
7. $y = c_1 + c_2 x + c_3 x^2 + \frac{7 \cos x}{9} - \frac{\cos^3 x}{27}.$

Art. 54.

1. $ax = \log(y + \sqrt{y^2 + c_1}) + c_2,$ or $y = c_1' e^{ax} + c_2' e^{-ax}.$
2. $ax = \sin^{-1} \frac{y}{c_1} + c_2,$ or $y = c_1 \sin(ax + c_2).$
3. $(c_1 x + c_2)^2 = a + c_1 y^2.$
4. $x\sqrt{2n} = c_1 \log \frac{\sqrt{c_1^2 e^{ny} + 1} - 1}{\sqrt{c_1^2 e^{ny} + 1} + 1} + c_2.$
5. $3x = 2a^{\frac{1}{3}}(y^{\frac{1}{2}} - 2c_1)(y^{\frac{1}{2}} + c_1)^{\frac{1}{2}} + c_2.$

Art. 56.

1. $y = c_1 \log x + c_2.$
2. $b^2 y = \log \sec[ab(x + c_1)] + c_2.$
3. $y = \frac{c_1 x^2}{2} + xf(c_1) + c_2.$
4. $\frac{2y}{a} = c_1 e^{\frac{x}{a}} + c_1^{-1} e^{-\frac{x}{a}} + c_2.$
5. $(x + c_1)^2 + (y + c_2)^2 = a^2.$
6. $y = c_1 x + (c_1^2 + 1) \log(x - c_1) + c_2.$
7. $y = c_1 \sin^{-1} x + (\sin^{-1} x)^2 + c_2.$
8. $y = \frac{4}{15c_1} (x + c_1^2 a^2)^{\frac{5}{2}} + c_2 x + c_3.$

9. $12y = w^3 + c_1w - 6w \log w + c_2$, where $w = x + c_3$.
 10. $y \sqrt{c_1} = w \sqrt{w^2 + c_1^2} + c_1^2 \log(w + \sqrt{w^2 + c_1^2}) + c_2$,
 where $w = x + c_3$.

Art. 58.

1. $y^2 = x^2 + c_1x + c_2$. 4. $c_1x = \log[c_1y + f(c_1)] + c_2$.
 2. $\log y - 1 = \frac{1}{c_1x + c_2}$. 5. $\log y = c_1e^x + c_2e^{-x}$.
 3. $c_1y = c_2e^{c_1x} - \sqrt{1 + a^2c_1^2}$. 6. $y^{n+1} = c_1e^{c_2x} + c_3$.

Art. 60.

1. $\begin{cases} 2x = (2c_2 - c_1 - c_2t)e^{-\frac{7t}{2}}, \\ y = (c_1 + c_2t)e^{-\frac{7t}{2}}. \end{cases}$
 2. $\begin{cases} y = -2c_1e^{-x} + c_2e^{-7x}, \\ z = c_1e^{-x} + c_2e^{-7x}. \end{cases}$
 3. $\begin{cases} 2x = e^{-6t}[(c_1 + c_2)\sin t + (c_2 - c_1)\cos t], \\ y = e^{-6t}(c_1\sin t + c_2\cos t). \end{cases}$
 4. $\begin{cases} y = c_1e^{nx} + c_2e^{-nx} + \frac{e^x}{n^2 - 1}, \\ az = -nc_1e^{nx} + nc_2e^{-nx} - \frac{e^x}{n^2 - 1}. \end{cases}$
 5. $\begin{cases} x = (c_1 + c_2t)e^{-4t} - \frac{e^{2t}}{36} + \frac{4e^t}{25}, \\ y = -(c_1 + c_2 + c_2t)e^{-4t} + \frac{7e^{2t}}{36} + \frac{e^t}{25}. \end{cases}$
 6. $\begin{cases} x = 2c_1e^{-4t} - c_2e^{-7t} + \frac{e^{2t}}{27} + \frac{7e^t}{40}, \\ y = c_1e^{-4t} + c_2e^{-7t} + \frac{7e^{2t}}{54} + \frac{e^t}{40}. \end{cases}$

$$7. \begin{cases} x = c_1 e^{-t} + c_2 e^{-6t} - \frac{29}{7} e^t + \frac{19}{3} t - \frac{56}{9}, \\ y = -c_1 e^{-t} + 4c_2 e^{-6t} + \frac{24}{7} e^t - \frac{17}{3} t + \frac{55}{9}. \end{cases}$$

$$8. \begin{cases} x = (c_1 + c_2 t) e^{-4t} - \frac{49}{36} e^{2t} + \frac{31}{25} e^t, \\ y = -(c_1 + c_2 + c_2 t) e^{-4t} + \frac{19}{36} e^{2t} - \frac{11}{25} e^t. \end{cases}$$

$$9. \begin{cases} x = (c_1 \sin t + c_2 \cos t) e^{-4t} + \frac{31}{26} e^t - \frac{93}{17}, \\ y = [(c_2 - c_1) \sin t - (c_2 + c_1) \cos t] e^{-4t} - \frac{2}{13} e^t + \frac{6}{17}. \end{cases}$$

$$10. \begin{cases} x = c_1 t^{-4} + 2c_2 t^{-3} + \frac{3}{10} t + \frac{t^2}{15}, \\ y = -c_1 t^{-4} - c_2 t^{-3} - \frac{t}{20} + \frac{2t^2}{15}. \end{cases}$$

$$11. \begin{cases} x = c_1 t^{-2} + \frac{t}{3}, \\ y = c_2 e^t - c_1 t^{-2} - \frac{t}{3}. \end{cases}$$

$$12. \begin{cases} x = a_1 \sin kt + a_2 \cos kt + a_3, \\ y = b_1 \sin kt + b_2 \cos kt + b_3, \\ z = c_1 \sin kt + c_2 \cos kt + c_3, \end{cases}$$

where $k^2 = l^2 + m^2 + n^2$.

The arbitrary constants are connected by the following equations :

$$\frac{mc_1 - nb_1}{a_2} = \frac{na_1 - lc_1}{b_2} = \frac{lb_1 - ma_1}{c_2} = k,$$

$$la_1 + mb_1 + nc_1 = 0, \quad \frac{a_3}{l} = \frac{b_3}{m} = \frac{c_3}{n}$$

$$13. \begin{cases} x = a_1 \sin(k \log t) + a_2 \cos(k \log t) + a_3, \\ y = b_1 \sin(k \log t) + b_2 \cos(k \log t) + b_3, \\ z = c_1 \sin(k \log t) + c_2 \cos(k \log t) + c_3, \end{cases}$$

where $k^2 = l^2 + m^2 + n^2$.

The arbitrary constants are connected by the following equations :

$$\frac{mn(c_1 - b_1)}{la_2} = \frac{nl(a_1 - c_1)}{mb_2} = \frac{lm(b_1 - a_1)}{nc_2} = k,$$

$$l^2 a_1 + m^2 b_1 + n^2 c_1 = 0, \quad a_3 = b_3 = c_3.$$

Art. 62.

1. $\begin{cases} x = c_1 \sin nt + c_2 \cos nt, \\ y = c_3 + c_4 t - x. \end{cases}$
2. $\begin{cases} x = (c_1 + c_2 t) e^t + (c_3 + c_4 t) e^{-t} - 23, \\ -2y = (c_1 - c_2 + c_2 t) e^t + (c_3 + c_4 + c_4 t) e^{-t} - 36. \end{cases}$
3. $\begin{cases} x = 4c_1 e^{2t} + 4c_2 e^{-2t} + c_3 e^{t\sqrt{7}} + c_4 e^{-t\sqrt{7}} + \frac{1}{7}, \\ y = c_1 e^{2t} + c_2 e^{-2t} + c_3 e^{t\sqrt{7}} + c_4 e^{-t\sqrt{7}} + \frac{9}{14}. \end{cases}$
4. $\begin{cases} x = e^{\frac{nt}{\sqrt{2}}} \left(c_1 \sin \frac{nt}{\sqrt{2}} + c_2 \cos \frac{nt}{\sqrt{2}} \right) + e^{-\frac{nt}{\sqrt{2}}} \left(c_3 \sin \frac{nt}{\sqrt{2}} + c_4 \cos \frac{nt}{\sqrt{2}} \right), \\ y = e^{\frac{nt}{\sqrt{2}}} \left(c_2 \sin \frac{nt}{\sqrt{2}} - c_1 \cos \frac{nt}{\sqrt{2}} \right) + e^{-\frac{nt}{\sqrt{2}}} \left(-c_4 \sin \frac{nt}{\sqrt{2}} + c_3 \cos \frac{nt}{\sqrt{2}} \right). \end{cases}$
5. $\begin{cases} y = (c_1 + c_2 x) e^x + 3c_3 e^{-\frac{3x}{2}} - \frac{x}{2}, \\ z = 2(3c_2 - c_1 - c_2 x) e^x - c_3 e^{-\frac{3x}{2}} - \frac{1}{3}. \end{cases}$
6. $y = u + v, \quad z = -u + v,$
 where $u = c_1 e^{2x} + (c_2 e^{nx} + c_3 e^{-nx}) e^x,$
 $v = c_4 e^{-2x} + (c_5 e^{nx} + c_6 e^{-nx}) e^{-x}.$

7. $y = u + v, \quad z = u - v,$

where $u = (c_1 + c_2x + c_3e^{x\sqrt{2}} + c_4e^{-x\sqrt{2}})e^x,$
 $v = (c_5 + c_6x + c_7e^{x\sqrt{2}} + c_8e^{-x\sqrt{2}})e^{-x}.$

Art. 63.

1. $x = cy^n.$

2. $y^2 = 4ax + c,$ a parabola.

3. $\pm(x+c) = \log(y + \sqrt{y^2-1}),$
 or $y = \frac{1}{2}(e^{x+c} + e^{-x-c}),$ a catenary.

4. $4my + c = 2mx\sqrt{4m^2x^2-1} + \log(2mx - \sqrt{4m^2x^2-1}).$

5. $\pm(x+c) = a \log(y + \sqrt{y^2-a^2}),$
 or $\pm(x+c) = a \log \frac{y + \sqrt{y^2-a^2}}{a},$
 from which $y = \frac{a}{2} \left(e^{\frac{x+c}{a}} + e^{-\frac{x+c}{a}} \right),$ a catenary.

6. $x^2 + y^2 = c^2,$ a circle.

7. $x^2 + y^2 - 2cy = 0,$ a circle.

8. $2x^2 + y^2 = 2c^2,$ an ellipse.

9. $x^2 - y^2 = c^2,$ an equilateral hyperbola.

10. $y^2 + x^2 = a^2 \log x^2 + c.$

11. $y = cx^m.$

12. $\frac{x^2}{h^2 - c^2} - \frac{y^2}{c^2} = 1,$ an ellipse or hyperbola.

13. $x^{\frac{1}{2}} + y^{\frac{1}{2}} = a^{\frac{1}{2}},$ a parabola.

14. $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}},$ a hypocycloid.

15. $2xy = a^2,$ an equilateral hyperbola.

16. $c^2y^2 - \log y^2 = 4c(x+c').$

17. $cy^2 - \frac{c^2}{n}(x+c')^2 = 1$, a hyperbola, when $n > 0$; an ellipse or hyperbola, when $n < 0$.

18. First, $(x+c')^2 + y^2 = c^2$, a circle.

Second, $\pm(x+c') = c \log(y + \sqrt{y^2 - c^2})$,

or $\pm(x+c') = c \log \frac{y + \sqrt{y^2 - c^2}}{c}$,

from which $y = \frac{c}{2} \left(e^{\frac{x+c'}{c}} + e^{-\frac{x+c'}{c}} \right)$, a catenary.

19. First, $x+c' = c \operatorname{vers}^{-1} \frac{y}{c} - \sqrt{2cy - y^2}$, a cycloid.

Second, $(x+c')^2 = 2cy - c^2$, a parabola.

ERRATA.



- p. 9, Ex. 12, for $= x^{a-1}dx$ read $= -x^{a-1}dx$.
- p. 11, Ex. 5, for $2y \sec x$ read $2y \sec y$.
- p. 21, Ex. 5, for $1 + (\log a)^2$ read $[1 + (\log a)^2]y$.
- p. 36, Art. 11, Ex. 7, for $2 \tan^{-1} \frac{y}{x}$ read $2 \tan^{-1} \frac{x}{y}$.
- p. 40, Art. 32, Ex. 7, for $\frac{\pm \sqrt{x^2+y} - x(\sqrt{17}-1)}{\pm \sqrt{x^2+y} + x(\sqrt{17}+1)}$
 read $\frac{\pm 4 \sqrt{x^2+y} - x(\sqrt{17}-1)}{\pm 4 \sqrt{x^2+y} + x(\sqrt{17}+1)}$.
- p. 40, Art. 34, Ex. 3, for $(y+c)^2 = \frac{(x+a)^2}{x^2+2ax}$
 read $e^{\frac{2y}{a}} + 2ce^{\frac{y}{a}}(x+a) + a^2c^2 = 0$.
- p. 41, Art. 42, Ex. 6, for $4c$ read $-4c$.
- p. 42, Art. 45, Ex. 14, for $\frac{1}{\sqrt{2}}$ and $-\frac{1}{\sqrt{2}}$
 read $\frac{x}{\sqrt{2}}$ and $-\frac{x}{\sqrt{2}}$.
- p. 44, Art. 48, Ex. 13, erase $-\frac{x}{4}$ and $+\frac{3}{8}$.
- p. 44, Art. 48, Ex. 14, erase $+\frac{33}{400}$ and $-\frac{3}{80}$.
- p. 44, Art. 50, Ex. 6, for $\frac{c_3+c_4 \log(x+1)}{\sqrt{x+1}}$
 read $\frac{c_3+c_4 \log(x+1)}{\sqrt{x+1}} + \frac{x^2+52x+51}{225}$.
- p. 45, Art. 54, Ex. 3, for a read $-a$.
- p. 46, Art. 56, Ex. 10, for $y\sqrt{c_1}$ read $2y\sqrt{c_1}$.
- p. 48, Art. 62, Ex. 4, in second members, for x read t .

A Treatise on Plane Surveying.

By DANIEL CARHART, C.E., Professor of Civil Engineering in the Western University of Pennsylvania, Allegheny. Illustrated. 8vo. Half leather. xvii + 498 pages. Mailing Price, \$2.00; for introduction, \$1.80.

THIS work covers the whole ground of Plane Surveying. It illustrates and describes the instruments employed, their adjustments and uses; it exemplifies the best methods of solving the ordinary problems occurring in practice, and furnishes solutions for many special cases which not infrequently present themselves. It is the result of twenty years' experience in the field and technical schools, and the aim has been to make it extremely practical, having in mind always that to become a *reliable* surveyor the student needs frequently to manipulate the various surveying instruments in the field, to solve many examples in the class-room, and to exercise good judgment in all these operations. Not only, therefore, are the different methods of surveying treated, and directions for using the instruments given, but these are supplemented by various field exercises to be performed, by numerous examples to be wrought, and by many queries to be answered.

Chapter I. Chain Surveying.

" II. Compass and Transit Surveying.

" III. Declination of the Needle.

" IV. Laying Out and Dividing Land.

" V. Plane Table Surveying.

" VI. Government Surveying.

" VII. City Surveying. Including the Principles of Levelling.

" VIII. Mine Surveying. Including the Elements of Topography.

The following **Tables** have been added: Logarithms of numbers; Approximate equation of time; Logarithms of trigonometric functions; For determining with greater accuracy than the preceding; Lengths of degrees of latitude and longitude; Miscellaneous formulas, and equivalents of metres, chains, and feet; Traverse; Natural sines and cosines; Natural tangents and cotangents.

The judicial functions of surveyors, as given by Chief Justice Cooley, are set forth in an appendix.

As a practical and complete treatise, Carhart's *Surveying* has received a cordial welcome.

A Short Table of Integrals.

Revised and Enlarged Edition. To accompany *Byerly's Integral Calculus*. By B. O. PEIRCE, Jr., Instructor in Mathematics, Harvard University. 32 pages. Mailing Price, 15 cents. Bound also with the Calculus.

Byerly's Syllabi.

By W. E. BYERLY, Professor of Mathematics in Harvard University. Each, 8 or 12 pages, 10 cents. The series includes, —

Syllabus of a Course in Plane Trigonometry.

Syllabus of a Course in Plane Analytical Geometry.

Syllabus of a Course in Plane Analytic Geometry. (Advanced Course.)

Syllabus of a Course in Analytical Geometry of Three Dimensions.

Syllabus of a Course on Modern Methods in Analytic Geometry.

Syllabus of a Course in the Theory of Equations.

Elements of the Differential and Integral Calculus.

With Examples and Applications. By J. M. TAYLOR, Professor of Mathematics in Madison University. 8vo. Cloth. 249 pages. Mailing Price, \$1.95; Introduction Price, \$1.80.

THE aim of this treatise is to present simply and concisely the fundamental problems of the Calculus, their solution, and more common applications.

Many theorems are proved both by the method of rates and that of limits, and thus each is made to throw light upon the other. The chapter on differentiation is followed by one on direct integration and its more important applications. Throughout the work there are numerous practical problems in Geometry and Mechanics, which serve to exhibit the power and use of the science, and to excite and keep alive the interest of the student.

The Nation, New York: It has two marked characteristics. In the first place, it is evidently a most carefully written book. . . . We are acquainted with no text-book of the calculus which compresses so much matter into so few pages, and at the same time leaves the impression that

all that is necessary has been said. In the second place, the number of carefully selected examples, both of those worked out in full in illustration of the text, and of those left for the student to work out for himself, is extraordinary.

Elementary Co-ordinate Geometry.

By W. B. SMITH, Professor of Physics, Missouri State University. 12mo. Cloth. 312 pages. Mailing Price, \$2.15; for introduction, \$2.00.

WHILE in the study of Analytic Geometry either gain of knowledge or culture of mind may be sought, the latter object alone can justify placing it in a college curriculum. Yet the subject may be so pursued as to be of no great educational value. Mere calculation, or the solution of problems by algebraic processes, is a very inferior discipline of reason. Even geometry is not the best discipline. In all thinking, the real difficulty lies in forming clear notions of things. In doing this all the higher faculties are brought into play. It is this formation of concepts, therefore, that is the essential part of mental training. And it is in line with this idea that the present treatise has been composed. Professors of mathematics speak of it as the most exhaustive work on the subject yet issued in America; and in colleges where an easier text-book is required for the regular course, this will be found of great value for post-graduate study.

Wm. G. Peck, *Prof. of Mathematics and Astronomy, Columbia College*: I have read Dr. Smith's Co-ordinate Geometry from beginning to end with unflagging interest. Its well compacted pages contain an immense amount of matter, most ad-

mirably arranged. It is an excellent book, and the author is entitled to the thanks of every lover of mathematical science for this valuable contribution to its literature. I shall recommend its adoption as a text-book in our graduate course.

Elements of the Theory of the Newtonian Potential Function.

By B. O. PEIRCE, Assistant Professor of Mathematics and Physics, Harvard University. 12mo. Cloth. 154 pages. Mailing Price, \$1.60; for introduction, \$1.50.

THIS book was written for the use of Electrical Engineers and students of Mathematical Physics because there was in English no mathematical treatment of the Theory of the Newtonian Potential Function in sufficiently simple form. It gives as briefly as is consistent with clearness so much of that theory as is needed before the study of standard works on Physics can be taken up with advantage. In the second edition a brief treatment of Electrokinematics and a large number of problems have been added.

Academic Trigonometry : Plane and Spherical.

By T. M. BLAKSLÉE, Ph.D. (Yale), Professor of Mathematics in the University of Des Moines. 12mo. Paper. 33 pages. Mailing Price, 20 cents; for introduction, 15 cents.

THE Plane and Spherical portions are arranged on opposite pages.

The memory is aided by analogies, and it is believed that the entire subject can be mastered in less time than is usually given to Plane Trigonometry alone, as the work contains but 29 pages of text.

The Plane portion is compact, and complete in itself.

Examples of Differential Equations.

By GEORGE A. OSBORNE, Professor of Mathematics in the Massachusetts Institute of Technology, Boston. 12mo. Cloth. vii + 50 pages. Mailing Price, 60 cents; for introduction, 50 cents.

A SERIES of nearly three hundred examples with answers, systematically arranged and grouped under the different cases, and accompanied by concise rules for the solution of each case.

Selden J. Coffin, *lately Prof. of Mathematics, Lafayette College* : Its appearance is most timely, and it supplies a manifest want.

Determinants.

The Theory of Determinants: an Elementary Treatise. By PAUL H. HANUS, B.S., recently Professor of Mathematics in the University of Colorado, now Principal of West High School, Denver, Col. 8vo. Cloth. viii + 217 pages. Mailing Price, \$1.90; for introduction, \$1.80.

THIS book is written especially for those who have had no previous knowledge of the subject, and is therefore adapted to self-instruction as well as to the needs of the class-room. The subject is at first presented in a very simple manner. As the reader advances, less and less attention is given to details. Throughout the entire work it is the constant aim to arouse and enliven the reader's interest, by first showing how the various concepts have arisen naturally, and by giving such applications as can be presented without exceeding the limits of the treatise.

William G. Peck, *Prof. of Mathematics, Columbia College, N.Y.* : A hasty glance convinces me that it is an improvement on Muir. (Aug. 30, 1886.)

T. W. Wright, *Prof. of Mathematics, Union Univ., Schenectady, N.Y.* : It fills admirably a vacancy in our mathematical literature, and is a very welcome addition indeed.

Analytic Geometry.

By A. S. HARDY, Ph.D., Professor of Mathematics in Dartmouth College, and author of *Elements of Quaternions*. 8vo. Cloth. xiv + 239 pages. Mailing Price, \$1.60; for introduction, \$1.50.

THIS work is designed for the student, not for the teacher.

Particular attention has been given to those fundamental conceptions and processes which, in the author's experience, have been found to be sources of difficulty to the student in acquiring a grasp of the subject as a *method of research*. The limits of the work are fixed by the time usually devoted to Analytic Geometry in our college courses by those who are not to make a special study in mathematics. It is hoped that it will prove to be a *text-book* which the teacher will wish to use in his class-room, rather than a *book of reference* to be placed on his study shelf.

Oren Root, *Professor of Mathematics, Hamilton College*: It meets quite fully my notion of a text for our classes. I have hesitated somewhat about introducing a generalized discussion of the conic in required work. I have, however, read Mr. Hardy's discussion carefully twice; and it seems to me that a student who can get the subject at all can get that. It is my present purpose to use the work next year.

John E. Clark, *Professor of Mathematics, Sheffield Scientific School of Yale College*: I need not hesitate to say, after even a cursory examination, that it seems to me a very attractive book, as I anticipated it

would be. It has evidently been prepared with real insight alike into the nature of the subject and the difficulties of beginners, and a very thoughtful regard to both; and I think its aims and characteristic features will meet with high approval. While leading the student to the usual useful results, the author happily takes especial pains to acquaint him with the character and spirit of analytical methods, and, so far as practicable, to help him acquire skill in using them.

John R. French, *Dean of College of Liberal Arts, Syracuse University*: It is a very excellent work, and well adapted to use in the recitation room.

Elements of Quaternions.

By A. S. HARDY, Ph.D., Professor of Mathematics, Dartmouth College. *Second edition, revised*. Crown 8vo. Cloth. viii + 234 pages. Mailing Price, \$2.15; Introduction, \$2.00.

THE chief aim has been to meet the wants of beginners in the class-room, and it is believed that this work will be found superior in fitness for beginners in practical compass, in explanations and applications, and in adaptation to the methods of instruction common in this country.

Elements of Plane Analytic Geometry.

By JOHN D. RUNKLE, Walker Professor of Mathematics in the Massachusetts Institute of Technology, Boston. 8vo. Cloth. ii + 344 pages. Mailing Price, \$2.25; for introduction, \$2.00.

IN this work, the author has had particularly in mind the needs of those students who can devote but a limited time to the subject, and yet must become quite familiar with at least its more elementary and fundamental part. For this reason, the earlier chapters are treated with somewhat more fulness than is usual. For some propositions, more than a single proof is given, and particular care has been taken to illustrate and enforce all parts of the subject by a large number of numerical applications. In the matter of problems, only the simpler ones have been selected, and the number has in every case been proportioned to the time that the students will have to devote to them. In general, propositions have been proved first with reference to rectangular axes. The determinant notation has not been used.

Descriptive Geometry.

By LINUS FAUNCE, Assistant Professor of Descriptive Geometry and Drawing in the Massachusetts Institute of Technology. 8vo. Cloth. 54 pages, with 16 lithographic plates, including 88 diagrams. Mailing Price, \$1.35; for introduction, \$1.25.

IN addition to the ordinary problems of Descriptive Geometry, this work includes a number of practical problems, such as might be met with by the draughtsman at any time, showing the application of the principles of Descriptive Geometry, a feature hitherto omitted in text-books on this subject. All of the problems have been treated clearly and concisely. The author's sole aim has been to present a work of practical value, not only as a text-book for schools and colleges, but also for every draughtsman.

The contents are: Chap. I., Elementary Principles; Notation. Chap. II., Problems relating to the Point, Line, and Plane. Chap. III., Principles and Problems relating to the Cylinder, Cone, and Double Curved Surfaces of Revolution. Chap. IV., Intersection of Planes and Solids, and the Development of Solids; Cylinders; Cones; Double Curved Surfaces of Revolution; Solids bounded by Plane Surfaces. Chap. V., Intersection of Solids. Chap. VI. Miscellaneous Problems.

CORNELL UNIVERSITY LIBRARY

OCT 20 1993

MATHEMATICS LIBRARY

